

# Amoxicillin plus clavulanic acid versus appendicectomy for treatment of acute uncomplicated appendicitis: an open-label, non-inferiority, randomised controlled trial



Corinne Vons, Caroline Barry\*, Sophie Maitre\*, Karine Pautrat, Mahaut Leconte, Bruno Costaglioli, Mehdi Karoui, Arnaud Alves, Bertrand Dousset, Patrice Valleur, Bruno Falissard, Dominique Franco

## Summary

**Background** Researchers have suggested that antibiotics could cure acute appendicitis. We assessed the efficacy of amoxicillin plus clavulanic acid by comparison with emergency appendicectomy for treatment of patients with uncomplicated acute appendicitis.

**Methods** In this open-label, non-inferiority, randomised trial, adult patients (aged 18–68 years) with uncomplicated acute appendicitis, as assessed by CT scan, were enrolled at six university hospitals in France. A computer-generated randomisation sequence was used to allocate patients randomly in a 1:1 ratio to receive amoxicillin plus clavulanic acid (3 g per day) for 8–15 days or emergency appendicectomy. The primary endpoint was occurrence of postintervention peritonitis within 30 days of treatment initiation. Non-inferiority was shown if the upper limit of the two-sided 95% CI for the difference in rates was lower than 10 percentage points. Both intention-to-treat and per-protocol analyses were done. This trial is registered with ClinicalTrials.gov, number NCT00135603.

**Findings** Of 243 patients randomised, 123 were allocated to the antibiotic group and 120 to the appendicectomy group. Four were excluded from analysis because of early dropout before receiving the intervention, leaving 239 (antibiotic group, 120; appendicectomy group, 119) patients for intention-to-treat analysis. 30-day postintervention peritonitis was significantly more frequent in the antibiotic group (8%, n=9) than in the appendicectomy group (2%, n=2; treatment difference 5·8; 95% CI 0·3–12·1). In the appendicectomy group, despite CT-scan assessment, 21 (18%) of 119 patients were unexpectedly identified at surgery to have complicated appendicitis with peritonitis. In the antibiotic group, 14 (12% [7·1–18·6]) of 120 underwent an appendicectomy during the first 30 days and 30 (29% [21·4–38·9]) of 102 underwent appendicectomy between 1 month and 1 year, 26 of whom had acute appendicitis (recurrence rate 26%; 18·0–34·7).

**Interpretation** Amoxicillin plus clavulanic acid was not non-inferior to emergency appendicectomy for treatment of acute appendicitis. Identification of predictive markers on CT scans might enable improved targeting of antibiotic treatment.

**Funding** French Ministry of Health, Programme Hospitalier de Recherche Clinique 2002.

## Introduction

Acute appendicitis is still the most common indication for surgery in patients admitted to hospital for acute abdominal pain. In about 20% of cases, acute appendicitis is complicated, leading to local or diffuse peritonitis;<sup>1</sup> most, however, are uncomplicated. Although urgent appendicectomy is still the recommended treatment for acute uncomplicated appendicitis, several studies, including four randomised trials,<sup>2–5</sup> have suggested that antibiotic treatment can cure acute appendicitis or can be the first line of treatment. Design limitations of previous studies, however, have decreased the relevance of their results, and consequently the current strategy for treatment of acute appendicitis has not been altered.<sup>6</sup> Although emergency appendicectomy is well tolerated by most patients, it is nevertheless associated with a risk of postoperative complications in about 2–23% of patients.<sup>7,8</sup> Additionally, over 10 years, 3% of patients

undergoing appendicectomy were readmitted for intestinal obstruction directly related to postoperative adhesions.<sup>9,10</sup> Avoidance of emergency appendicectomy in patients with uncomplicated appendicitis, who otherwise would have had surgery, would therefore improve the risk–benefit ratio of acute-appendicitis treatment. We compared the results of treatment with amoxicillin plus clavulanic acid with emergency appendicectomy in a group of patients with uncomplicated acute appendicitis as assessed by CT.

## Methods

### Patients

We undertook an open-label, non-inferiority, randomised controlled trial. The study took place in six academic centres of the Assistance Publique-Hôpitaux de Paris, France, and was approved by the ethics committee of the Hôpital Bicêtre, Le Kremlin-Bicêtre, France. All patients provided signed, informed consent.

*Lancet* 2011; 377: 1573–79

See [Comment](#) page 1545

\*These authors contributed equally

Hôpital Antoine Bécélère (Assistance Publique-Hôpitaux de Paris and Université Paris XI), Service de Chirurgie

(Prof C Vons MD, Prof D Franco MD), et de Radiologie (S Maitre MD), Clamart, France; Inserm, U669, Université Paris-Sud and Université Paris Descartes, Paris,

France AP-HP, Hôpital Paul Brousse, Villejuif, France (C Barry PhD, Prof B Falissard MD); Hôpital Lariboisière (Assistance

Publique-Hôpitaux de Paris and Université Paris VII), Service de Chirurgie, Paris, France


(K Pautrat MD, Prof A Alves MD, Prof P Valleur MD); Hôpital Cochin (Assistance Publique-Hôpitaux de Paris and Université Paris V), Service de Chirurgie, Paris, France (M Leconte MD,

Prof B Dousset MD); Hôpital Bicêtre (Assistance Publique-Hôpitaux de Paris and Université Paris XI), Service de Chirurgie, Paris, France (B Costaglioli MD); and

Le Kremlin Bicêtre and Hôpital Henri Mondor (Assistance Publique-Hôpitaux de Paris and Université Paris XII), Service de Chirurgie, Créteil, France (M Karoui MD)

Correspondence to: Prof Corinne Vons, Service de Chirurgie, Hôpital Jean Verdier, Avenue du 14 Juillet, Bondy 93143, France [corinne.vons@jvr.aphp.fr](mailto:corinne.vons@jvr.aphp.fr)

For the trial protocol see [http://www.medicine.u-psud.fr/fr/recherche/les\\_publications/trialappendicitis.html](http://www.medicine.u-psud.fr/fr/recherche/les_publications/trialappendicitis.html)



**Figure: Trial profile**

ITT=intention to treat. \*Withdrew consent before starting treatment.

All adults examined in the emergency department and suspected to have an acute appendicitis were assessed for possible inclusion in the study. Patients were excluded if one of the following criteria were present: age less than 18 years (no upper age limit); antibiotic treatment 5 days before; allergy to  $\beta$ -lactam antibiotics; known intolerance to amoxicillin plus clavulanic acid (nausea, vomiting); receiving steroid or anticoagulant treatments; past history of inflammatory bowel disease; pregnancy or a positive pregnancy test; life expectancy less than 1 year; allergy to iodine or blood creatinine 200  $\mu$ mol/L or more; or inability to understand information about the protocol or to sign the consent form.

Patients eligible for inclusion to the study were informed of the protocol and invited to participate. After informed consent was obtained, a CT scan was done. Diagnosis of uncomplicated appendicitis was assessed by CT imaging. An emergency radiologist in the hospital

where the patient was admitted did a CT scan of patients' appendices according to the standard protocol of the hospitals. CT imaging was done with a 16-multidetector CT scanner in all centres, except for one (Hôpital Antoine Bécélère), which used a single-row-detector spiral CT scanner (webappendix p 1).

A final diagnosis of uncomplicated acute appendicitis required clear visualisation of the appendix (appendix diameter >6 mm and no opacification of the appendix in patients with enema), and absence of any of the three following criteria of complicated appendicitis with peritonitis: extra luminal gas, periappendiceal fluid, or disseminated intraperitoneal fluid. An appendix diameter greater than 15 mm was a criterion for exclusion from the study, because of risk of malignancy.<sup>11</sup> Caecal-wall thickening, inflammation of periappendiceal fat, and presence of intraluminal stercoliths were also recorded, but were not exclusion criteria.

See Online for webappendix

### Randomisation and masking

When a diagnosis of uncomplicated acute appendicitis was made, patients were individually assigned to undergo either appendectomy or treatment with amoxicillin plus clavulanic acid. The computer-generated randomisation code was produced by the trial statistician. To ensure balance between the numbers in each group, block sizes of four were generated for allocation of patients to one of the two treatment groups; the randomisation procedure was stratified by site, with an equal allocation ratio. Opaque, sealed, and sequentially numbered envelopes were provided to each trial site. To enrol a patient, an independent pharmacologist opened the next consecutively numbered envelope.

### Procedures

Patients were admitted to hospital irrespective of the treatment assigned and were assessed twice a day while in hospital. They were discharged after resolution of pain, fever, and any digestive symptoms. Appendectomy was done according to surgeons' standard practice (a McBurney incision or laparoscopy). Patients were given one injection of amoxicillin plus clavulanic acid (2 g) at induction of general anaesthesia, but did not receive antibiotic treatment thereafter, unless complicated appendicitis was diagnosed during surgery, in which case patients were given postoperative antibiotics.

Patients in the antibiotic treatment group received amoxicillin plus clavulanic acid (3 g per day for patients weighing <90 kg, and 4 g per day for patients ≥90 kg), given intravenously to those with nausea or vomiting, and orally to all others.<sup>12-14</sup> This drug combination was chosen because of its efficacy for ambulatory treatment of uncomplicated sigmoiditis.<sup>13</sup> If symptoms and abdominal tenderness did not resolve after 48 h, immediate appendectomy was undertaken. If pain and fever resolved rapidly, patients were discharged. Patients continued the same antibiotic treatment at home, with the same dose, for 8 days, and were seen on day 8; persistence of pain or fever prompted a CT scan and possible appendectomy. In the absence of these symptoms, a sustained high white-blood-cell count or a high C-reactive-protein concentration resulted in extension of antibiotic treatment for a further 8 days. Persistence of similar biological disorders on day 15 prompted appendectomy without an additional CT scan. All patients were seen systematically in consultations on days 15, 30, 90, 180, and 360.

Histological examination of the appendix was done after every appendectomy. Definitive diagnosis of uncomplicated acute appendicitis was based on the presence of mucosal ulceration with neutrophil infiltration restricted to the mucosa, or with a transmural extension. The primary binary endpoint was occurrence of peritonitis within 30 days of initial treatment. In the antibiotic group, diagnosis of peritonitis was done either by appendectomy when a complicated appendicitis was identified, or postoperatively by CT scan. In the appendectomy group,

|   | Appendectomy group (n=119) | Antibiotic treatment group (n=120) |
|---|----------------------------|------------------------------------|
| Age (years)                             | 34 (12)                    | 31 (9)                             |
| Sex                                     |                            |                                    |
| Men                                     | 70 (59%)                   | 73 (61%)                           |
| Women                                   | 49 (41%)                   | 47 (39%)                           |
| Body-mass index (kg/m <sup>2</sup> )    | 24.1 (4.1)                 | 23.0 (3.0)                         |
| Employment status                       |                            |                                    |
| Full-time or part-time work             | 87 (73%)                   | 83 (69%)                           |
| Clinical symptoms on admission          |                            |                                    |
| Sudden onset of pain*                   | 59 (50%)                   | 57 (48%)                           |
| Mean pain score VAS (0-10)†             | 6.4 (2.1)                  | 6.3 (1.9)                          |
| Body temperature >37.5°C                | 32 (28%)                   | 38 (32%)                           |
| Lower-right quadrant-rebound tenderness | 72 (62%)                   | 56 (48%)                           |
| Biological findings                     |                            |                                    |
| Leucocytes (10 <sup>9</sup> /L)         | 13.1 (3.4)                 | 13.6 (3.6)                         |
| High CRP concentration‡                 | 78 (68%)                   | 76 (68%)                           |
| Additional CT findings§                 |                            |                                    |
| Local caecal-wall thickening            | 14 (13%)                   | 17 (15%)                           |
| Inflammation of periappendiceal fat     | 47 (44%)                   | 49 (44%)                           |
| Intraluminal stercolith                 | 22 (21%)                   | 19 (17%)                           |

Data are mean (SD) or numbers (%). We calculated percentages using non-missing data. VAS=visual analogue scale. CRP=C-reactive protein. \*All patients had abdominal pain, but only a proportion presented with sudden onset pain. †Intensity of pain was measured with a horizontal 10-mm visual analogue scale, with 0 representing no pain, and 10 the worst pain ever experienced. ‡High CRP denotes concentrations above normal values as defined by hospital laboratories; definition of high CRP was beyond a specific value; normal values differed between hospitals. §CT findings additional to inclusion and exclusion criteria (appendix diameter >6 mm and no extra luminal gas, periappendiceal fluid, and disseminated intraperitoneal fluid).

**Table 1: Baseline characteristics of the intention-to-treat population**

|   | Appendectomy group (n=119) | Antibiotic-treatment group (n=120) | Difference (95% CI)   |
|---|----------------------------|------------------------------------|-----------------------|
| <b>Primary endpoint events</b>                                  |                            |                                    |                       |
| 30-day post-therapeutic peritonitis                             | 2 (2%)*                    | 9 (8%)†                            | 5.8 (0.3 to 12.1)     |
| <b>Incidence of peritonitis</b>                                 |                            |                                    |                       |
| Complicated appendicitis with peritonitis identified at surgery | 21 (18%)‡                  | 9 (8%)†                            | -10.1 (-18.7 to -1.7) |
| Postoperative peritonitis                                       | 2 (2%)‡                    | 2 (2%)§                            | 0 (-4.4 to 4.4)       |

Data are number unless otherwise stated. \*In the appendectomy group, two cases of postoperative peritonitis occurred; these patients had postoperative localised peritonitis successfully treated with antibiotics. †In the antibiotic group, complicated appendicitis with peritonitis was identified during appendectomy performed within 30 days of treatment initiation in nine of 14 patients who did not show improvement. ‡Discovery of a complicated appendicitis with peritonitis in the appendectomy group was not a primary endpoint. §Two patients in the antibiotic group, who underwent secondary appendectomy, had postoperative peritonitis.

**Table 2: Incidence of primary endpoint events and complicated appendicitis with peritonitis and postoperative peritonitis within 30 days after the start of treatment in both groups (intention-to-treat population)**

diagnosis of postoperative peritonitis was made with CT scan findings for patients with fever, abdominal pain, and high concentrations of white-blood cells and C reactive protein. Signs of localised postoperative peritonitis on CT scans were densification of soft tissue with or without organised fluid collection (abscess) of the right iliac fossa. Appendectomy done within 30 days of treatment

|                           | Appendicectomy group (n=119) | Antibiotic-treatment group (n=120) | p value |
|---------------------------|------------------------------|------------------------------------|---------|
| Duration of pain*         | 1.70 (1.07)                  | 1.63 (1.35)                        | 0.13    |
| Duration of hospital stay | 3.04 (1.50)                  | 3.96 (4.87)                        | 0.08    |
| Duration of disability    | 10.45 (8.20)                 | 9.82 (10.51)                       | 0.25    |

Data are mean (SD), p value (Wilcoxon test). Duration in days. \*Pain as assessed with the visual analogue score  $\geq 4$ .

**Table 3: Duration of post-therapeutic pain, hospital stay, and disability (intention-to-treat population)**

|                                | Within 30 days (n=120) | Between 30 days and 1 year of follow-up (n=102)* |
|--------------------------------|------------------------|--|
| Number of patients (%; 95% CI) | 14 (12%; 7.1–18.6)     | 30 (29%; 21.4–38.9)                              |
| Appendicitis (%; 95% CI)       | 13 (11%; 6.4–17.7)     | 26 (25%; 18.0–34.7)                              |
| Complicated†                   | 9                      | 3  |
| Uncomplicated                  | 4                      | 23   |
| No appendicitis                | 1                      | 4  |
| Fibrous                        | 1                      | 4  |

\*120 patients in the antibiotic-treatment group minus 14 patients who had an appendicectomy during the first 30 days of follow-up and four who discontinued follow-up before 1 month. †Complicated appendicitis with peritonitis.

**Table 4: Aspects of appendices during appendicectomies done in 44 of 120 patients treated initially with antibiotics (intention-to-treat population)**

initiation in the antibiotic group was not a primary endpoint if complicated appendicitis with peritonitis was not identified at surgery.

Secondary endpoints were the number of days with a postintervention visual-analogue-scale pain score  $\geq 4$  (on a 0–10 scale),<sup>15</sup> length of hospital stay and absence from work (total days including any additional hospital stays), incidence of complications other than peritonitis within 1 year (postoperative wound abscess, incisional hernia, adhesive occlusion), and recurrence of appendicitis after antibiotic treatment (appendicectomy done between 30 days and 1 year of follow-up, with a confirmed diagnosis of appendicitis).

### Statistical analysis

This study was based on the notion that antibiotic treatment would not be inferior to appendicectomy in relation to the primary efficacy outcome, with the use of a prespecified non-inferiority margin—the upper limit of the 95% CI for the difference in rates would not exceed 10 percentage points.

We calculated that a sample size of 200 patients would give a power of 80% to establish whether antibiotic treatment was not inferior to appendicectomy in relation to the 30-day incidence of postintervention peritonitis. This sample size took into account an expected 30-day incidence of peritonitis after appendicectomy for

uncomplicated appendicitis of 2%,<sup>16,17</sup> a non-inferiority margin of 10%, and a two-sided  $\alpha$  risk of 0.05.<sup>18</sup> However, we planned to enrol 250 patients because of the possible loss of patients after their inclusion.

Study outcomes were assessed by both intention-to-treat and per-protocol analyses. The intention-to-treat population included all randomised participants who began a treatment (surgical treatment or at least one dose of antibiotics). A second reading of the CT scan was done later by an assigned non-emergency radiologist, who was masked to the patients' treatment or status, to confirm initial CT scan diagnosis of uncomplicated appendicitis made by the emergency radiologist. The per-protocol population included all patients who completed the study (1 year), and for whom the second reading of a CT-scan confirmed the diagnosis of uncomplicated appendicitis.

The primary analysis in this non-inferiority trial compared the two study groups for the rate of peritonitis that occurred within 30 days of treatment initiation. The 95% CIs for absolute difference in percentages between the antibiotic-treatment group and the surgery group were estimated according to the methodology used by Altman and colleagues.<sup>19</sup> Secondary binary endpoints were similarly analysed. Wilcoxon tests were used to compare durations. Webappendix pp 1–2 shows management of missing data. All reported p values are two-sided and were not adjusted for multiple testing.

We did additional post-hoc analyses. Factors predictive of complicated appendicitis in the appendicectomy group were calculated. For the antibiotic-treatment group, factors predictive of absence of improvement (patients needing appendicectomy during the first 30 days after start of antibiotic treatment, with a confirmed histological diagnosis of acute appendicitis), or of recurrence of appendicitis (patients needing appendicectomy between 30 days and 1 year, with a confirmed histological diagnosis of acute appendicitis) were also calculated. Univariate logistic-regression models were used to assess the association between these events and each patient's baseline clinical characteristics. We used R software (version 2.7.0) for all analyses.

### Role of the funding source

The sponsor of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. CV, CB, BF, and DF had full access to all data in the study and final responsibility for the decision to submit for publication. All other authors could request examination of any of the data elements.

### Results

The figure shows the trial profile. 243 patients (aged 18–68 years) were enrolled into the study between March 11, 2004, and January 16, 2007. Four refused to participate in the trial shortly after randomisation, therefore 239 patients constituted the intention-to-treat population. Table 1 shows baseline characteristics of these patients.

Table 2 shows incidence of primary endpoints and incidence of complicated appendicitis with peritonitis and postoperative peritonitis within 30 days after the start of treatment in the intention-to-treat population. 30 day postintervention peritonitis was significantly more frequent in the antibiotic group than in the appendectomy group.

24 (10%) of 239 patients did not complete the 1-year study. 11 were excluded by re-reading of the CT-scan. The remaining 204 patients constituted the per-protocol population (figure; results are shown in webappendix pp 2–5).

Primary endpoint data were missing for nine patients (four in the antibiotic group and five in the appendectomy group). We did a sensitivity analysis to establish the effect of missing cases, excluding patients with missing data from the analysis. The rate of peritonitis within 30 days of treatment initiation remained higher in the antibiotic-treatment group than in the surgery group (difference 6.0 percentage points; 95% CI 0.3–12.5; webappendix pp 5–6).

For secondary endpoints, the median duration of severe pain, days in hospital, and absence from work did not differ between the two groups (table 3). Other postintervention complications included postoperative wound infection (two of 120 in the antibiotic group vs one of 119 in the appendectomy group) and intestinal adhesive occlusion (one of 120 in the antibiotic group vs none in the appendectomy group) during the 1-year follow-up. Incisional hernia did not occur in either group. No significant differences were identified between the two groups for any postintervention complications.

Table 4 shows the aspects of the appendices assessed in patients who had appendectomies treated initially with antibiotics in the intention-to-treat population. Those who underwent appendectomy between 1 month and 1 year had the operation after a median of 4.2 months (range 1.2–11.1). Overall, 81 (68%) of 120 patients did not need an appendectomy for acute appendicitis in the antibiotic group during the 1-year follow-up.

Post-hoc analyses showed that laparoscopic or McBurney approach rates were similar in both groups (webappendix pp 6–7). Logistic-regression analyses showed that CT-scanner type (multidetector vs single detector) was not significantly associated with misdiagnosed complicated appendicitis ( $p=0.42$ ; table 5). Presence of a stercolith on preoperative CT scan was the only factor associated with a significantly increased risk of complicated appendicitis (table 5,  $p<0.0001$ ). In the antibiotic group, the presence of a stercolith was also the only factor associated with failure of antibiotic treatment for appendicitis (table 6,  $p=0.0072$ ).

In the subgroup of patients without visualisation of a stercolith on initial CT scan, we identified no significant difference in the incidence of 30-day postintervention peritonitis between the two groups (difference 2.9 percentage points; 95% CI –3.0 to 9.2;

|   | Complicated appendicitis (n=21) | Uncomplicated appendicitis (n=98) |
|---|---------------------------------|-----------------------------------|
| Age (years)                             | 38 (13)                         | 34 (12)                           |
| Sex                                     |                                 |                                   |
| Men                                     | 16 (76%)                        | 54 (55%)                          |
| Women                                   | 5 (24%)                         | 44 (45%)                          |
| Clinical symptoms on admission          |                                 |                                   |
| Sudden onset of pain*                   | 11 (52%)                        | 48 (48%)                          |
| Mean pain score (0–10)†                 | 6.3 (1.7)                       | 6.4 (2.2)                         |
| Body temperature (>37°C)                | 9 (43%)                         | 23 (24%)                          |
| Lower-right quadrant-rebound tenderness | 13 (62%)                        | 59 (61%)                          |
| Biological findings                     |                                 |                                   |
| Leucocyte counts ( $10^9/L$ )           | 13.9 (2.6)                      | 13.0 (3.6)                        |
| High CRP concentration‡                 | 16 (80%)                        | 62 (66%)                          |
| Additional appendix CT findings§        |                                 |                                   |
| Local caecal-wall thickening            | 2 (12%)                         | 12 (13%)                          |
| Inflammation of periappendiceal fat     | 11 (65%)                        | 36 (40%)                          |
| Intraluminal stercolith                 | 9 (53%)                         | 13 (15%)                          |
| CT-scanner type                         |                                 |                                   |
| Multidetector                           | 17 (81%)                        | 71 (72%)                          |

Data are mean (SD) or numbers (%). Comparison of the 21 cases of complicated appendicitis with peritonitis versus 98 cases of uncomplicated appendicitis or no appendicitis, identified at surgery. CRP=C-reactive protein. \*All patients had abdominal pain, but only some presented with sudden onset pain. †Pain assessed with the visual analogue score >4. ‡High C-reactive protein denotes concentrations above normal values defined by hospital laboratories; definition of high CRP was beyond a specific value; normal values differed between hospitals. §CT findings additional to inclusion and exclusion criteria (appendix diameter >6 mm and no extra luminal gas, periappendiceal fluid, and disseminated intraperitoneal fluid).

**Table 5: Association between risk factors and complicated appendicitis with peritonitis in the appendectomy group**

webappendix p 7). No factors were associated with the recurrence of appendicitis. No adverse events were deemed by the investigator as being related to CT-scanning or antibiotic treatment.

## Discussion

Incidence of 30-day postintervention peritonitis, which was the main judgment criterion, was significantly higher in the antibiotic-treatment group than in the appendectomy group. This study showed that antibiotic treatment with amoxicillin plus clavulanic acid was not non-inferior to emergency appendectomy for treatment of acute uncomplicated appendicitis.

Trials that show that acute appendicitis can be treated successfully with antibiotics<sup>2–5</sup> were weakened by several design limitations. For example, diagnosis of uncomplicated appendicitis was not supported by systematic CT-scan assessment, although researchers claimed to have treated uncomplicated appendicitis alone.<sup>3</sup> Therefore, we tried to avoid these limitations in our study by using CT scans to select patients with uncomplicated appendicitis before randomisation. Multiple detector CT scanning is generally accepted as the best investigation to diagnose acute appendicitis, because it has a high sensitivity and specificity.<sup>20</sup> Indeed, only 3% of patients allocated to surgery in our trial had

|   | No improvement of appendicitis (n=13) | Without appendectomy during the first month (n=102) |
|---|---------------------------------------|---|
| Age (years)                             | 35 (13)                               | 31 (9)  |
| Sex                                     |                                       |   |
| Men                                     | 10 (77%)                              | 59 (58%)  |
| Women                                   | 3 (23%)                               | 43 (42%)  |
| Clinical symptoms on admission          |                                       |   |
| Sudden onset of pain*                   | 8 (62%)                               | 47 (46%)  |
| Mean pain score (0–10)†                 | 6.4 (1.9)                             | 6.2 (2.0)   |
| Body temperature (>37°C)                | 6 (46%)                               | 29 (29%)  |
| Lower-right quadrant-rebound tenderness | 7 (54%)                               | 46 (46%)  |
| Biological findings                     |                                       |   |
| Leucocyte counts (10 <sup>9</sup> /L)   | 13.9 (3.3)                            | 13.5 (3.7)  |
| High CRP concentration‡                 | 9 (90%)                               | 64 (67%)  |
| Additional appendix CT findings§        |                                       |   |
| Local caecal-wall thickening            | 1 (8%)                                | 15 (16%)  |
| Inflammation of periappendiceal fat     | 8 (67%)                               | 40 (43%)  |
| Intraluminal stercolith                 | 6 (50%)                               | 13 (14%)  |

Data are means (SD) or numbers (%). We calculated percentages using non-missing data. Comparison of 13 cases with absence of improvement after antibiotic treatment versus 102 without appendectomy during the first month. VAS=visual analogue scale. CRP=C-reactive protein. \*All patients had abdominal pain, but only some presented with sudden onset pain. †Pain assessed with the visual analogue score >4. ‡High C reactive protein denotes concentrations above normal values as defined by hospital laboratories. §CT findings additional to inclusion and exclusion criteria (appendix diameter >6 mm and no extra luminal gas, periappendiceal fluid, and disseminated intraperitoneal fluid).

**Table 6: Association between risk factors and absence of improvement of appendicitis in the antibiotic-treatment group**

no appendicitis, which compares favourably with the 10–15% reported in two previous studies.<sup>4,5</sup>

The study was limited by the short follow-up period. Recurrence of appendicitis might have continued after one year. Masking of participants or clinicians to treatment allocation was not possible, and research assessors were also not masked.

In our trial, two-thirds of patients in the antibiotic group who needed an appendectomy during the 30 days after treatment initiation had complicated appendicitis, consistent with previous studies.<sup>3,4</sup> This finding could be interpreted as a failure of the antibiotics to prevent complications after non-operated acute uncomplicated appendicitis; however, if this were the case, the rate of complicated appendicitis discovered during appendectomy in the antibiotic group would be expected to be higher than that identified in the appendectomy group. In fact, complicated appendicitis was less frequent in the antibiotic group than in the appendectomy group (table 2). Alternatively, complicated appendicitis might already have been present in these patients at the time of randomisation, despite not being diagnosed on CT scan, and some were successfully treated with amoxicillin plus clavulanic acid.

Therefore, our finding that antibiotic treatment with amoxicillin plus clavulanic acid was inferior relative to appendectomy in patients with uncomplicated acute appendicitis might be related to the small proportion of

### Panel: Research in context

#### Systematic review

Four trials comparing antibiotic treatment with emergency appendectomy have been reported.<sup>2-5</sup> They all concluded that acute appendicitis can be treated successfully with antibiotics,<sup>2-5</sup> and some suggested that antibiotics might be used as first-line treatment for unselected patients with acute appendicitis.<sup>4</sup> A meta-analysis of three trials showed a trend towards a lower risk of complications in the antibiotic-treated group than in the surgical group.<sup>6</sup> However, results of these trials had several protocol design limitations, especially the loose criteria chosen for the definition of acute appendicitis—diagnosis of acute appendicitis was based on clinical examination,<sup>2,3</sup> the clinical-biological score,<sup>5</sup> and ultrasonography of the right iliac fossa,<sup>4</sup> but a CT scan before randomisation was not used to confirm the diagnosis. Because of absence of morphological investigation in the surgical group of two large trials, 10–15% of patients did not have appendicitis,<sup>4,5</sup> and 39–48% had complicated appendicitis.<sup>3,4</sup>

High rate of protocol violation was also a limitation in one study—50% of patients allocated to antibiotic treatment underwent appendectomy immediately after randomisation.<sup>4</sup> Rate of patients lost to follow-up by 1 year in two trials were problematic:<sup>3,4</sup> in one the rate was very high<sup>4</sup> and in the other it was not clearly reported.<sup>3</sup> Therefore, results from these studies might underestimate the true recurrence rate. Consequently, whether antibiotics are a genuinely valid alternative to appendectomy for treatment of acute uncomplicated appendicitis remains unclear.

#### Interpretation

Our trial is the first trial in which patients were included and randomised patients only after proven diagnosis of acute uncomplicated appendicitis by CT scanning. Treatment with amoxicillin and clavulanic acid was not as effective as emergency appendectomy for treatment of acute appendicitis. Our results show that exploration of predictive markers on CT scans might allow improvement in patient selection for antibiotic-based treatment.

patients with complicated appendicitis who were erroneously included and randomised. Distinction between uncomplicated and complicated appendicitis remains difficult even with multiple-detector CT scans.<sup>21,22</sup> Morphological diagnosis of appendiceal perforation depends on indirect but late signs, which are very specific but have a low sensitivity; however, Tsuboi and colleagues<sup>23</sup> have suggested that morphological diagnosis could be improved. Additionally, in our trial and other reports, visualisation of a stercolith on the initial CT scan predicted both complicated appendicitis in patients treated with appendectomy<sup>22</sup> and failure in the antibiotic group.<sup>24</sup> Even though complicated appendicitis can also be cured with antibiotics, further trials of such treatment of acute

appendicitis should focus on use of new diagnostic techniques for improved patient selection.

The inferiority of antibiotic treatment versus appendectomy could be also related to appendicitis resistant to amoxicillin plus clavulanic acid. Evidence shows that resistance of *Escherichia coli* to this antibiotic combination is increasing.<sup>25</sup> Third-generation cephalosporins could be used, although they are not yet recommended.<sup>26</sup>

Nearly a quarter of our patients who recovered after antibiotic treatment had recurrence of appendicitis; this finding is more than the 14% reported in previous studies.<sup>3,4</sup> This difference could be explained by the high rate of patients lost to 1-year follow-up in previous trials.<sup>3,4</sup> Our results suggest that emergency appendectomy remains the gold standard for treatment of acute uncomplicated appendicitis (panel).

#### Contributors

CV conceived, designed, implemented, and led the study. CV, SM, KP, ML, BC, MK, AA, BD, and PV contributed to the recruitment, clinical care, and follow-up of patients. CV, CB, and DF analysed and managed data. SM re-read all CT scans. BF designed the plan for statistical analysis, and CB did the statistical analysis. CV, CB, BF, and DF drafted the report, contributed to the interpretation of results, and prepared the report. All authors reviewed and approved the final version of the manuscript.

#### Conflicts of interest

We declare that we have no conflicts of interest.

#### Acknowledgments

We thank A Moschetto for help with identification of patients lost to follow-up; Solene Fabre for her contribution to data management; Laurent Becquemont for his constant support; and D Poussin, R Balzarotti, A Carloni, J M Proske, I Dagher, S Gueroult, J M Thillois, P Taleb, S Servadjian, J L Bouillot, I Lorand, F Gayral, A Laurent, and D Cherqui for their contributions to the study. This trial was supported by a grant from the Projet Hospitalier de Recherche Clinique of the French Ministry of Health (AOR 02 063) and Assistance Publique-Hôpitaux de Paris.

#### References

- Livingston EH, Woodward WA, Sarosi GA, Haley RW. Disconnect between incidence of nonperforated and perforated appendicitis: implications for pathophysiology and management. *Ann Surg* 2007; **245**: 886–92.
- Eriksson S, Granstrom L. Randomized controlled trial of appendectomy versus antibiotic therapy for acute appendicitis. *Br J Surg* 1995; **82**: 166–69.
- Styrud J, Eriksson S, Nilsson I, et al. Appendectomy versus antibiotic treatment in acute appendicitis: a prospective multicenter randomized controlled trial. *World J Surg* 2006; **30**: 1033–37.
- Hansson J, Korner U, Khorram-Manesh A, Solberg A, Lundholm K. Randomized clinical trial of antibiotic therapy versus appendectomy as primary treatment of acute appendicitis in unselected patients. *Br J Surg* 2009; **96**: 473–81.
- Malik AA, Bari SU. Conservative management of acute appendicitis. *J Gastrointest Surg* 2009; **13**: 966–70.
- Varadhan KK, Humes DJ, Neal KR, Lobo DN. Antibiotic therapy versus appendectomy for acute appendicitis: a meta-analysis. *World J Surg* 2010; **34**: 199–209.
- Konstantinidis KM, Anastasakou KA, Vorias MN, Sambalis GH, Georgiou MK, Xiarchos AG. A decade of laparoscopic appendectomy: presentation of 1,026 patients with suspected appendicitis treated in a single surgical department. *J Laparoendosc Adv Surg Tech A* 2008; **18**: 248–58.
- Ming PC, Yan TY, Tat LH. Risk factors of postoperative infections in adults with complicated appendicitis. *Surg Laparosc Endosc Percutan Tech* 2009; **19**: 244–48.
- Parker MC, Ellis H, Moran BJ, et al. Postoperative adhesions: ten-year follow-up of 12,584 patients undergoing lower abdominal surgery. *Dis Colon Rectum* 2001; **44**: 822–29.
- Leung TT, Dixon E, Gill M, et al. Bowel obstruction following appendectomy: what is the true incidence? *Ann Surg* 2009; **250**: 51–53.
- Pickhardt PJ, Levy AD, Rohrmann CA Jr, Kende AI. Primary neoplasms of the appendix manifesting as acute appendicitis: CT findings with pathologic comparison. *Radiology* 2002; **224**: 775–81.
- Pelaez N, Pera M, Courtier R, et al. Applicability, safety and efficacy of an ambulatory treatment protocol in patients with uncomplicated acute diverticulitis. *Cir Esp* 2006; **80**: 369–72.
- Schug-Pass C, Geers P, Hugel O, Lippert H, Kockerling F. Prospective randomized trial comparing short-term antibiotic therapy versus standard therapy for acute uncomplicated sigmoid diverticulitis. *Int J Colorectal Dis* 2010; **25**: 751–59.
- Fraser JD, Aguayo P, Leys CM, et al. A complete course of intravenous antibiotics vs a combination of intravenous and oral antibiotics for perforated appendicitis in children: a prospective, randomized trial. *J Pediatr Surg* 2009; **45**: 1198–202.
- Lingreen R, Grider JS. Retrospective review of patient self-reported improvement and post-procedure findings for mild (minimally invasive lumbar decompression). *Pain Physician* 2010; **13**: 555–60.
- Aguilo J, Peiro S, Munoz C, et al. Adverse outcomes in the surgical treatment of acute appendicitis. *Cir Esp* 2005; **78**: 312–17.
- Ingraham AM, Cohen ME, Bilimoria KY, Pritts T1A, Ko CY, Esposito TJ. Comparison of outcomes after laparoscopic versus open appendectomy for acute appendicitis at 222 ACS NSQIP hospitals. *Surgery* 2010; **148**: 625–35.
- Piaggio G, Elbourne DR, Altman DG, Pocock SJ, Evans SJ. Reporting of noninferiority and equivalence randomized trials: an extension of the CONSORT statement. *JAMA* 2006; **295**: 1152–60.
- Altman DG, Gore SM, Gardner MJ, Pocock SJ. Statistical guidelines for contributors to medical journals. In: Altman DG, Machin D, Bryant TN, Gardner MJ, eds. *Statistics with Confidence: Confidence Intervals and Statistical Guidelines*. 2nd edn. London: BMJ Books; 2000: 171–90.
- Horton MD, Counter SF, Florence MG, Hart MJ. A prospective trial of computed tomography and ultrasonography for diagnosing appendicitis in the atypical patient. *Am J Surg* 2000; **179**: 379–81.
- Pinto Leite N, Pereira JM, Cunha R, Pinto P, Sirlin C. CT evaluation of appendicitis and its complications: imaging techniques and key diagnostic findings. *Am J Roentgenol* 2005; **185**: 406–17.
- Bixby SD, Lucey BC, Soto JA, Theysohn JM, Ozonoff A, Varghese JC. Perforated versus nonperforated acute appendicitis: accuracy of multidetector CT detection. *Radiology* 2006; **241**: 780–86.
- Tsuboi M, Takase K, Kaneda I, et al. Perforated and nonperforated appendicitis: defect in enhancing appendiceal wall: depiction with multi-detector row CT. *Radiology* 2008; **246**: 142–47.
- Shindoh J, Niwa H, Kawai K, et al. Predictive factors for negative outcomes in initial non-operative management of suspected appendicitis. *J Gastrointest Surg* 2009; **14**: 309–14.
- Pitout JDD, Laupland KB. Extended-spectrum  $\beta$ -lactamase-producing *Enterobacteriaceae*: an emerging public-health concern. *Lancet Infect Dis* 2008; **8**: 159–66.
- de Kraker ME, Wolkewitz M, Davey PG et al. Burden of antimicrobial resistance in European hospitals: excess mortality and length of hospital stay associated with bloodstream infections due to *Escherichia coli* resistant to third-generation cephalosporins. *J Antimicrob Chemother* 2010; **66**: 398–407.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.